

Poster presentation

Relationship of various body composition parameters used for predicting VO_{2max}

Julie Culbertson*¹, Allyn Byars², Susan Keith² and Mike Greenwood¹

Address: ¹Baylor University, Waco, 00000, TX, USA and ²Angelo State University, San Angelo, TX 76909, USA

Email: Julie Culbertson* - Julie_Culbertson@baylor.edu

* Corresponding author

from 2008 International Society of Sports Nutrition Conference and Expo Las Vegas, NV, USA. 9–10 June 2008

Published: 17 September 2008

Journal of the International Society of Sports Nutrition 2008, **5**(Suppl 1):P12 doi:10.1186/1550-2783-5-S1-P12

This abstract is available from: <http://www.jissn.com/content/5/S1/P12>

© 2008 Culbertson et al; licensee BioMed Central Ltd.

Background

Forty-six male and twenty-eight female university students ($n = 74$), ages 18–32 (21.41 ± 2.54) volunteered to participate in a study examining the relationship of various body parameters used for predicting aerobic power (VO_{2max}).

Methods

Participants performed a graded exercise treadmill test measuring VO_{2max} as well as a 6-minute walk test within 1 week of each other. Height, weight, self-report of physical activity (PARS), body mass index (BMI), and estimated percent body fat (skinfold and bioelectrical impedance analysis) were also measured including the recording of participant demographics.

Results

Using multiple linear regression, a model was produced where significant predictors of VO_{2max} ($p < 0.05$) included estimated body fat percentage from sum of three skinfolds ($p < .001$) and BMI ($p = .024$). The generalized equation ($R = .741$, $R^2 = .549$, $SEE = 7.10$) derived from the model to predict VO_{2max} was: $72.32 + (-.949 \times \%fat \text{ from skinfolds}) - (.392 \times BMI)$. Variables excluded ($p > 0.05$) from the model included 6-minute walk test time, PARS, height, weight, gender, age, and percent body fat from bioelectrical impedance analysis.

Conclusion

It was concluded that that percent body fat and BMI are significant predictors of VO_{2max} . Although the relationship between VO_{2max} and the other variables examined in this study were not significant, it may have been due to the homogeneity of these parameters (e.g., age and activity levels) in the participants and the lack of the 6-minute walk test to discriminate among fitness levels in university students. However, these results demonstrate the importance of the relationship between BMI and percent body fat (estimated from skinfolds) in predicting VO_{2max} without maximal testing.